

New results on synthesis of Hs isotopes

J. Dvorak¹, R. Krücken¹, F. Nebel¹, Z. Novackova¹, A. Türler¹, B. Wierczinski¹, A. Yakushev¹, W. Brühle², E. Jäger², E. Schimpf², M. Schädel², A. Semchenkov^{1,2}, A. Kuznetsov³, M. Chelnokov³, A. Yeremin³, Ch. E. Düllmann⁴, Y. Nagame⁵, K. Eberhardt⁶, P. Thörle⁶, R. Dressler⁷, Z. Qin^{2,8}, M. Wegrzecki⁹

¹TUM Garching; ²GSI Darmstadt; ³JINR Dubna; ⁴LBNL Berkeley and UC Berkeley;
⁵JAERI Tokai; ⁶Universität Mainz; ⁷PSI Villigen; ⁸IMP Lanzhou, ⁹ITE Warsaw

Synthesis and investigation of heavy nuclei close to the deformed nuclear shells at $Z=108$ and $N=162$, especially of the even-even nucleus ^{270}Hs , provide important data for comparison with theoretical predictions. Relatively long half-lives for alpha decay and spontaneous fission were predicted for these nuclei [1,2]. The first chemical identification and the study of nuclear decay properties of Hs isotopes was accomplished in 2001 [3,4]. Gas phase chemical separation of Hs isotopes in the form of volatile HsO_4 provides a highly selective and very efficient method of isolation. Here we report preliminary results of a recent Hs chemistry experiment performed at the GSI UNILAC accelerator to measure the excitation function of the reaction $^{26}\text{Mg}(^{248}\text{Cm}, xn)^{274-xn}\text{Hs}$ and to obtain additional information on the decay properties of Hs isotopes and their daughters. Based on calculated excitation functions, experiments at beam energies of $E_{\text{lab}} = 135$ MeV and $E_{\text{lab}} = 145$ MeV in the center of the target were carried out. The resulting excitation energies of 40 MeV and 49 MeV correspond to the predicted maxima of the 4n and 5n evaporation channel, respectively. The experimental procedure is described in [5]. Beam

integrals of 1.46×10^{18} (145 MeV) and 2.02×10^{18} (135 MeV) ^{26}Mg ions were accumulated. A first, preliminary analysis of the data at 145 MeV revealed 7 decay chains attributed to the decay of Hs isotopes. These decay chains with observed α -particle or fission fragment energies in MeV, decay times, as well as detector number, top (T) or bottom (B) position and gas flow rates are presented in Figure 1. The probabilities that the observed decay chains are of random origin are low and comparable to the values given in [3,4]. The data accumulated at 135-MeV is currently being analyzed.

References

- [1] I. Muntian et al., Acta Phys. Pol. B **32**, 691 (2001).
- [2] R. Smolańczuk et al., Phys. Rev. C, **52**, 1871 (1995).
- [3] Ch.E. Düllmann et al., Nature **418**, 859 (2002).
- [4] A. Türler et al., Eur. Phys. J. A **17**, 505 (2003).
- [5] J. Dvorak et al., this Annual Report, p.




Figure 1. Correlated decay chains observed in the 145-MeV run.